Grass and Forage Science (1996) Volume 51, 466–468

Author Index

ANNICCHIARICO P. *see* PIANO E.
ANNICCHIARICO P., BOZZO F., PARENTE G., GUSMEROLI F., MAIR V., MARGUERETTAZ O. and ORLAND D. Analysis of adaptation of grass/legume mixtures to Italian alpine and subalpine zones through an additive main effects and multiplicative interaction model 405
ARMSTRONG R.H., ROBERTSON E. and HUNTER E.A. The effect of sward height and its direction of change on the herbage intake, diet selection and performance of weaned lambs grazing ryegrass swards 389

BAILEY G. *see* SMITH K.A.
BALFOURIER F. *see* RAVEL C.
BARTHRAM G.T. and GRANT S.A. Interactions between variety and the timing of conservation cuts on species balance in *Lolium Perenne-Trifolium repens* swards 98
BINNIE R.C. and FROST J.P. Some effects of applying undiluted silage effluent to grassland 272
BOZZO F. *see* ANNICCHIARICO P.
BULLARD M.J. and CRAWFORD T.J. Productivity of *Lotus corniculatus* L. (bird's-foot trefoil) in the UK when grown under low-input conditions as spaced plants, monoculture swards or mixed swards 439

CALVIÈRE I. and DURU M. Leaf appearance and senescence patterns of some pasture species 447
CHAMBERLAIN D.G. *see* WAYMAN J.
CHAMPION R.A. *see* ORR R.J.
CHARMET G. *see* RAVEL C.
CHERRILL A. Infestation of improved grasslands by *Juncus effusus* L. in the catchment of the River Tyne, Northern England: a field survey (Research Note) 85
COOPER A. *see* WELLER R.F.
CRAWFORD T.J. *see* BULLARD M.J.
CRESSWELL A. *see* MATTHEW C.
CRITTEN D.L. *see* WILLIAMS A.G.
CROSSE S. *see* DILLON P.
CUSSEN R.F., MERRY R.J., WILLIAMS A.P. and TWEED J.K.S. The effect of additives on the ensilage of forage of differing perennial ryegrass and white clover content 249

DANIEL G.J. *see* DAVIES D.A.
DAVIES D.A., FOTHERGILL M., DANIEL G.J. and MORGAN C.T. Animal production evaluation of herbage varieties. 2. Comparison of Aberystwyth S184, AberEndura and Grasslands Huia white clovers 227
DHANOA M.S. *see* MERRY R.J.
DHANOA M.S. *see* WELLER R.F.
DILLON P., CROSSE S., STAKELUM G. and FLYNN F. The effect of calving date and stocking rate on the performance of spring-calving dairy cows 286
DINDAL D.L. *see* MURPHY W.M.
DOWDESWELL A.M. *see* FISHER G.E.J.
DOWDESWELL A.M. *see* FISHER G.E.J.
DURU M. *see* CALVIÈRE I.

EDGAR K.F., KLESSA D.A., FRAME J. and HARKESS R.D. The nitrogen value of anaerobically digested sewage sludge applied to grassland during winter 353
EDWARDS G.R., PARSONS A.J., PENNING P.D. and NEWMAN J.A. Relationship between vegetation state and bite dimensions of sheep grazing contrasting plant species and its implications for intake rate and diet selection 378
EVANS D.R., WILLIAMS T.A., JONES S. and EVANS S.A. The effect of blending white clover varieties and their contribution to a mixed grass/clover sward under continuous sheep stocking 10
EVANS S.A. *see* EVANS D.R.

FEIST D. *see* HAMILTON W.J.
FISHER A. and WILMAN D. Effect of interval between harvests and spring-applied fertilizer N on the growth of white clover in a mixed sward 162
FISHER G.E.J. and DOWDESWELL A.M. The performance of summer-calving cows grazing perennial ryegrass swards 315
FISHER G.E.J. and ROBERTS D.J. The effects of stocking rate in spring, and topping on sward characteristics and dairy cow performance in poor grass growing season 461
FISHER G.E.J., ROBERTS D.J. and DOWDESWELL A.M. The manipulation of grass swards for summer-calving dairy cows 424
FLYNN F. *see* DILLON P.
FOTHERGILL M. *see* DAVIES D.A.
FRAME J. *see* EDGAR K.F.
FROST J.P. *see* BINNIE R.C.
FROST J.P., POOTS R., KNIGHT A., GORDON F.J. and LONG F.N.J. Effect of forage matting rate on grass drying, rate of silage fermentation, silage intake and silage digestibility, 21
FULKERSON W.J. and SLACK K. Leaf number as a criterion for determining defoliation time for *Lolium perenne*: 2. Effect of defoliation frequency and height 16

GHESQUIÈRE M. *see* HAZARD L.
GIBB M.J. *see* WILKINS R.J.
GIBSON G.J. *see* RUXTON G.D.
GORDON F.J. *see* FROST J.P.
GRANT S.A. *see* BARTHRAM G.T.
GUSMEROLI F. *see* ANNICCHIARICO P.

HAGGAR R.J. *see* MATTHEW C.
HAIGH P.M. The composition of first-cut grass for ensilage in England and Wales from 1988 to 1991 (Research Note) 63
HAMILTON W.J., SIBBALD M.A. and FEIST D. The effect of sward height on the liveweight gain of farmed yearling red deer (*Cervus elaphus*) stags 399
HANDA D.P., SREENATH P.R. and RAJPALI S.K. Uniformity trial with lucerne grown for fodder 209
HARKESS R.D. *see* EDGAR K.F.
HAYES M.J. and WILLIAMS E.D. Effects of timing and intensity of defoliation on growth and development of white clover seedlings 414
HAYNES R.J. *see* WILLIAMS P.H.
HAZARD L. and GHESQUIÈRE M. Evidence from the use of isozyme markers of competition in swards between short-leaved and long-leaved perennial ryegrass 241

© 1996 Blackwell Science Ltd.

Grass and Forage Science (1996) Volume 51, 466–468

Author Index

ANNICCHIARICO P. *see* PIANO E.
ANNICCHIARICO P., BOZZO F., PARENTE G., GUSMEROLI F., MAIR V., MARGUERETTAZ O. and ORLAND D. Analysis of adaptation of grass/legume mixtures to Italian alpine and subalpine zones through an additive main effects and multiplicative interaction model 405
ARMSTRONG R.H., ROBERTSON E. and HUNTER E.A. The effect of sward height and its direction of change on the herbage intake, diet selection and performance of weaned lambs grazing ryegrass swards 389

BAILEY G. *see* SMITH K.A.
BALFOURIER F. *see* RAVEL C.
BARTHRAM G.T. and GRANT S.A. Interactions between variety and the timing of conservation cuts on species balance in *Lolium Perenne-Trifolium repens* swards 98
BINNIE R.C. and FROST J.P. Some effects of applying undiluted silage effluent to grassland 272
BOZZO F. *see* ANNICCHIARICO P.
BULLARD M.J. and CRAWFORD T.J. Productivity of *Lotus corniculatus* L. (bird's-foot trefoil) in the UK when grown under low-input conditions as spaced plants, monoculture swards or mixed swards 439

CALVIÈRE I. and DURU M. Leaf appearance and senescence patterns of some pasture species 447
CHAMBERLAIN D.G. *see* WAYMAN J.
CHAMPION R.A. *see* ORR R.J.
CHARMET G. *see* RAVEL C.
CHERRILL A. Infestation of improved grasslands by *Juncus effusus* L. in the catchment of the River Tyne, Northern England: a field survey (Research Note) 85
COOPER A. *see* WELLER R.F.
CRAWFORD T.J. *see* BULLARD M.J.
CRESSWELL A. *see* MATTHEW C.
CRITTEN D.L. *see* WILLIAMS A.G.
CROSSE S. *see* DILLON P.
CUSSEN R.F., MERRY R.J., WILLIAMS A.P. and TWEED J.K.S. The effect of additives on the ensilage of forage of differing perennial ryegrass and white clover content 249

DANIEL G.J. *see* DAVIES D.A.
DAVIES D.A., FOTHERGILL M., DANIEL G.J. and MORGAN C.T. Animal production evaluation of herbage varieties. 2. Comparison of Aberystwyth S184, AberEndura and Grasslands Huia white clovers 227
DHANOA M.S. *see* MERRY R.J.
DHANOA M.S. *see* WELLER R.F.
DILLON P., CROSSE S., STAKELUM G. and FLYNN F. The effect of calving date and stocking rate on the performance of spring-calving dairy cows 286
DINDAL D.L. *see* MURPHY W.M.
DOWDESWELL A.M. *see* FISHER G.E.J.
DOWDESWELL A.M. *see* FISHER G.E.J.
DURU M. *see* CALVIÈRE I.

EDGAR K.F., KLESSA D.A., FRAME J. and HARKESS R.D. The nitrogen value of anaerobically digested sewage sludge applied to grassland during winter 353
EDWARDS G.R., PARSONS A.J., PENNING P.D. and NEWMAN J.A. Relationship between vegetation state and bite dimensions of sheep grazing contrasting plant species and its implications for intake rate and diet selection 378
EVANS D.R., WILLIAMS T.A., JONES S. and EVANS S.A. The effect of blending white clover varieties and their contribution to a mixed grass/clover sward under continuous sheep stocking 10
EVANS S.A. *see* EVANS D.R.

FEIST D. *see* HAMILTON W.J.
FISHER A. and WILMAN D. Effect of interval between harvests and spring-applied fertilizer N on the growth of white clover in a mixed sward 162
FISHER G.E.J. and DOWDESWELL A.M. The performance of summer-calving cows grazing perennial ryegrass swards 315
FISHER G.E.J. and ROBERTS D.J. The effects of stocking rate in spring, and topping on sward characteristics and dairy cow performance in poor grass growing season 461
FISHER G.E.J., ROBERTS D.J. and DOWDESWELL A.M. The manipulation of grass swards for summer-calving dairy cows 424
FLYNN F. *see* DILLON P.
FOTHERGILL M. *see* DAVIES D.A.
FRAME J. *see* EDGAR K.F.
FROST J.P. *see* BINNIE R.C.
FROST J.P., POOTS R., KNIGHT A., GORDON F.J. and LONG F.N.J. Effect of forage matting rate on grass drying, rate of silage fermentation, silage intake and silage digestibility, 21
FULKERSON W.J. and SLACK K. Leaf number as a criterion for determining defoliation time for *Lolium perenne*: 2. Effect of defoliation frequency and height 16

GHESQUIÈRE M. *see* HAZARD L.
GIBB M.J. *see* WILKINS R.J.
GIBSON G.J. *see* RUXTON G.D.
GORDON F.J. *see* FROST J.P.
GRANT S.A. *see* BARTHRAM G.T.
GUSMEROLI F. *see* ANNICCHIARICO P.

HAGGAR R.J. *see* MATTHEW C.
HAIGH P.M. The composition of first-cut grass for ensilage in England and Wales from 1988 to 1991 (Research Note) 63
HAMILTON W.J., SIBBALD M.A. and FEIST D. The effect of sward height on the liveweight gain of farmed yearling red deer (*Cervus elaphus*) stags 399
HANDA D.P., SREENATH P.R. and RAJPALI S.K. Uniformity trial with lucerne grown for fodder 209
HARKESS R.D. *see* EDGAR K.F.
HAYES M.J. and WILLIAMS E.D. Effects of timing and intensity of defoliation on growth and development of white clover seedlings 414
HAYNES R.J. *see* WILLIAMS P.H.
HAZARD L. and GHESQUIÈRE M. Evidence from the use of isozyme markers of competition in swards between short-leaved and long-leaved perennial ryegrass 241

© 1996 Blackwell Science Ltd.

HIGUCHI S. *see* PASUMARTY S.V.
HIGUCHI S. *see* PASUMARTY S.V.
HILL M.J., HOCKNEY M.J., MULCAHY C.A. and RAPP G. The effect of hydrogen cyanide potential (HCNp) and sward morphology on the relative acceptability to sheep of white clover and Caucasian clover herbage 1
HOCKNEY M.J. *see* HILL M.J.
HODGSON I. *see* SMITH K.A.
HUCKLE C.A. *see* WILKINS R.J.
HUNTER E.A. *see* ARMSTRONG R.H.
HUTCHINGS N.J. and KRISTENSEN I.S. Modelling mineral nitrogen accumulation in grazed pasture: will more nitrogen leach from fertilized grass than unfertilized grass/clover? 300

JACKSON D.R. *see* SMITH K.A.
JONES S. *see* EVANS D.R.

KAFFKA S.R. *see* KANNEGANTI V.R.
KANNEGANTI V.R. and KAFFKA S.R. Forage availability from a temperate pasture managed with intensive rotational grazing 55
KEADY T.W.J. and STEEN R.W.J. The effects of treating low dry-matter, low digestibility grass with a bacterial inoculant on the intake and performance of beef cattle, and studies on its mode of action 217
KIRKHAM F.W. and TALLOWIN J.R.B. The influence of cutting date and previous fertilizer treatment on the productivity and botanical composition of species-rich hay meadows on the Somerset Levels 365
KLESSA D.A. *see* EDGAR K.F.
KNIGHT A. *see* FROST J.P.
KRISTENSEN I.S. *see* HUTCHINGS N.J.

LAIDLAW A.S. *see* PATTERSON J.D.
LAIDLAW A.S., WITHERS J.A. and TOAL L.G. The effect of surface height of swards continuously stocked with cattle on herbage production and clover content over four years 48
LINDGREN S. *see* ÖSTLING C.
LONG F.N.J. *see* FROST J.P.

MAIR V. *see* ANNICCHIARICO P.
MARGUERETTAZ O. *see* ANNICCHIARICO P.
MARSHALL A.H. Peduncle characteristics, inflorescence survival and reproductive growth of white clover (*Trifolium repens* L.) 324
MATSUMARA T. *see* PASUMARTY S.V.
MATSUMARA T. *see* PASUMARTY S.V.
MATTHEW C., CRESSWELL A. and HAGGAR R.J. Characteristics of microsites with and without white clover in two field swards (Research Note) 178
MCBRIDE J. *see* PATTERSON J.D.
MENA BARRETO A.D. *see* MURPHY W.M.
MENA BARRETO A.D. *see* MURPHY W.M.
MENA BARRETO A.D. *see* MURPHY W.M.
MERCHANT M. The effect of pattern and severity of cutting on the vigour of the soft rush (*Juncus effusus* L.) (Research Note) 81
MERRY R.J. *see* CUSSEN R.F.
MERRY R.J., DHANOA M.S. and THEODOROU M.K. Use of freshly cultured lactic acid bacteria as silage inoculants 112
MORGAN C.T. *see* DAVIES D.A.
MORRIS R.M. *see* STUDDY C.D.
MULCAHY C.A. *see* HILL M.J.
MURATA T. *see* PASUMARTY S.V.
MURATA T. *see* PASUMARTY S.V.
MURPHY W.M., MENA BARRETO A.D. and SILMAN J.P. Sward dynamics of a smooth-stalked meadowgrass-dominant white clover sward rotationally grazed by cattle and/or sheep 183
MURPHY W.M., MENA BARRETO A.D., SILMAN J.P. and DINDAL D.L. Cattle and sheep grazing effects on soil organisms, fertility and compaction in a smooth-stalked meadowgrass-dominant white clover sward 191
MURPHY W.M., SILMAN J.P. and MENA BARRETO A.D. A comparison of quadrat, capacitance meter, HFRO sard stick, and rising plate for estimating herbage mass in a smooth-stalked, meadowgrass-dominant white clover sward 452

NEWMAN J.A. *see* EDWARDS G.R.

ORLAND D. *see* ANNICCHIARICO P.
ORR R.J., PENNING P.D., PARSONS A.J. and CHAMPION R.A. Herbage intake and N excretion by sheep grazing monocultures or a mixture of grass and white clover 31
ÖSTLING C. and LINDGREN S. Influences of enterobacteria on the fermentation and aerobic stability of grass silages 41

PAPASTYLIANOU I. Effect of rainfall and temperature on yield of *Vicia sativa* under rainfed Mediterranean conditions 456
PARENTE G. *see* ANNICCHIARICO P.
PARSONS A.J. *see* EDWARDS G.R.
PARSONS A.J. *see* ORR R.J.
PASUMARTY S.V., HIGUCHI S., MURATA T. and MATSUMARA T. Influence of seed quality on seedling growth of white clover (*Trifolium repens* L.) 93
PASUMARTY S.V., HIGUCHI S., MURATA T. and MATSUMARA T. Seed production in white clover (*Trifolium repens* L.). II. Effect of autumn defoliation on potential seed yield and seed yield components 147
PATTERSON J.D., LAIDLAW A.S. and MCBRIDE J. The influence of autumn management and companion grass on the development of white clover over winter in mixed swards 345
PENNING P.D. *see* EDWARDS G.R.
PENNING P.D. *see* ORR R.J.
PIANO E. and ANNICCHIARICO P. Persistance of Ladino white clover ecotypes and its relationship with other agronomic traits 195
POOTS R. *see* FROST J.P.

RAJPALI S.K. *see* HANDA D.P.
RAPP G. *see* HILL M.J.
RAVEL C., CHARMET G. and BALFOURIER F. Influence of the fungal endophyte *Acremonium lolii* on agronomic traits of perennial ryegrass in France (Research Note) 75
REYNOLDS A.M. *see* WILLIAMS A.G.
RIDGE I. *see* STUDDY C.D.
ROBERTS D.J. *see* FISHER G.E.J.
ROBERTS D.J. *see* FISHER G.E.J.
ROBERTSON E. *see* ARMSTRONG R.H.
RUXTON G.D. and GIBSON G.J. A mathematical model of aerobic deterioration of big-bale silage and its implications for the growth of *Listeria monocytogenes* 331

SIBBALD M.A. *see* HAMILTON W.J.
SILMAN J.P. *see* MURPHY W.M.
SILMAN J.P. *see* MURPHY W.M.
SILMAN J.P. *see* MURPHY W.M.
SIMONS R.G. A non-destructive method for estimating lucerne shoot weights (Research Note) 68
SLACK K. *see* FULKERSON W.J.
SMITH K.A., JACKSON D.R., UNWIN R.J., BAILEY G. and HODGSON I. Negative effects of winter- and spring-applied cattle slurry on the yield of herbage at simulated early grazing and first-cut silage 124
SREENATH P.R. *see* HANDA D.P.
STAKELUM G. *see* DILLON P.

STEEN R.W.J. *see* KEADY T.W.J.
STUDDY C.D., MORRIS R.M. and RIDGE I. The effects of separated cow slurry liquor on soil and herbage nitrogen in *Phalaris arundinacea* and *Lolium perenne* 106

TALLOWIN J.R.B. *see* KIRKHAM F.W.
THEODOROU M.K. *see* MERRY R.J. 259
THOMPSON L. Sites of photoperception in white clover
TOAL L.G. *see* LAIDLAW A.S.
TWEED J.K.S. *see* CUSSEN R.F.

UNWIN R.J. *see* SMITH K.A.

WAYMAN J., CHAMBERLAIN D.G. and WILLIAMS A.G. A comparison of the ensilage, in an anaerobic cabinet, of grass either fresh or when thawed after deep freezing (Research Note) 71
WELLER R.F., COOPER A. and DHANOA M.S. The selection of winter wheat varieties for whole crop cereal conservation (Research Note) 172
WILKINS P.W. Independence of dry matter yield and leaf yield among perennial ryegrass varieties differing in seasonal yield distribution 155
WILKINS R.J., GIBB M.J. and HUCKLE C.A. Lactation performance of spring-calving dairy cows grazing mixed perennial ryegrass/white clover swards of differing composition and height 199
WILLIAMS A.G. *see* WAYMAN J.
WILLIAMS A.G., CRITTEN D.L. and REYNOLDS A.M. A mathematical model of the aerobic deterioration of silage 132
WILLIAMS A.P. *see* CUSSEN R.F.
WILLIAMS E.D. *see* HAYES M.J.
WILLIAMS P.H. and HAYNES R.J. Effect of sheep, deer and cattle dung on herbage production and soil nutrient content 263
WILLIAMS T.A. *see* EVANS D.R.
WILMAN D. *see* FISHER A.
WITHERS J.A. *see* LAIDLAW A.S.